

Documentation Table 5

Live births by birthweight and race of mother and infant deaths and infant mortality rates by age at death, birthweight, and race of mother for 10 major causes of infant death: United States, 2000 birth cohort data

[Infant deaths are weighted. Infant deaths are under 1 year. Neonatal deaths are under 28 days; early neonatal, 0-6 days; late neonatal, 7-27 days; and postneonatal, 28 days through 11 months. Rates are per 100,000 live births]

Cause of death, birthweight, and race of mother	Live births	Infant deaths	Total neonatal	Early neonatal	Late neonatal	Post-neonatal
All races ¹						
All birthweights						
All causes	4,058,903	28,049	18,734	14,885	3,850	9,315
Rate		691.1	461.6	366.7	94.9	229.5
Congenital malformations (Q00-Q99)		5,769	4,163	3,183	980	1,606
Rate		142.1	102.6	78.4	24.1	39.6
Short gestation and low birthweight nec (P07)		4,404	4,323	4,196	127	81
Rate		108.5	106.5	103.4	3.1	2.0
Sudden infant death syndrome (R95)		2,485	201	30	171	2,284
Rate		61.2	5.0	0.7	4.2	56.3
Maternal complications of pregnancy (P01)		1,390	1,378	1,363	15	12
Rate		34.3	34.0	33.6	*	*
Complications of placenta, cord, membranes (P02) ..		1,044	1,028	990	37	16
Rate		25.7	25.3	24.4	0.9	*
Respiratory distress of newborn (P22)		1,003	931	746	185	72
Rate		24.7	22.9	18.4	4.6	1.8
Accidents (unintentional injuries) (V01-X59)		897	90	28	61	808
Rate		22.1	2.2	0.7	1.5	19.9
Bacterial sepsis of newborn (P36)		775	745	334	411	30
Rate		19.1	18.4	8.2	10.1	0.7
Diseases of the circulatory system (I00-I99)		680	251	139	112	429
Rate		16.7	6.2	3.4	2.8	10.6
Intrauterine hypoxia and birth asphyxia (P20-P21)		619	583	423	160	36
Rate		15.2	14.4	10.4	3.9	0.9
All other causes		8,983	5,043	3,452	1,590	3,941
Rate		221.3	124.2	85.1	39.2	97.1
Less than 2,500 grams						
All causes		18,218	14,923	12,527	2,397	3,295
Rate		5,913.6	4,844.0	4,066.0	777.9	1,069.6
Congenital malformations (Q00-Q99)		3,246	2,587	2,187	400	659
Rate		1,053.6	839.8	709.9	129.9	213.8
Short gestation and low birthweight nec (P07)		4,194	4,113	3,989	125	80
Rate		1,361.2	1,335.1	1,294.7	40.4	26.1
Sudden infant death syndrome (R95)		491	43	8	35	448
Rate		159.5	14.1	*	11.5	145.4
Maternal complications of pregnancy (P01)		1,306	1,295	1,281	14	11
Rate		423.9	420.3	415.7	*	*
Complications of placenta, cord, membranes (P02) ..		905	893	869	24	11
Rate		293.6	290.0	282.2	7.8	*
Respiratory distress of newborn (P22)		973	908	727	181	66
Rate		316.0	294.7	235.9	58.8	21.3
Accidents (unintentional injuries) (V01-X59)		126	12	8	4	114
Rate		41.0	*	*	*	37.0
Bacterial sepsis of newborn (P36)		654	631	277	354	23
Rate		212.2	204.7	89.9	114.8	7.5

See footnotes at end of table.

Documentation Table 5

Live births by birthweight and race of mother and infant deaths and infant mortality rates by age at death, birthweight, and race of mother for 10 major causes of infant death: United States, 2000 birth cohort data - Con.

[Infant deaths are weighted. Infant deaths are under 1 year. Neonatal deaths are under 28 days; early neonatal, 0-6 days; late neonatal, 7-27 days; and postneonatal, 28 days through 11 months. Rates are per 100,000 live births]

Cause of death, birthweight, and race of mother	Live births	Infant deaths	Total neonatal	Early neonatal	Late neonatal	Post-neonatal
All races ¹						
Less than 2,500 grams						
Diseases of the circulatory system (I00-I99)		280	117	72	44	163
Rate		90.9	37.9	23.5	14.4	53.1
Intrauterine hypoxia and birth asphyxia (P20-P21)		289	282	223	58	7
Rate		93.7	91.4	72.5	18.9	*
All other causes		5,755	4,042	2,885	1,157	1,713
Rate		1,868.0	1,312.1	936.6	375.5	555.9
2,500 grams or more						
All causes		9,424	3,434	1,987	1,447	5,990
Rate		251.4	91.6	53.0	38.6	159.8
Congenital malformations (Q00-Q99)		2,492	1,550	971	579	942
Rate		66.5	41.3	25.9	15.4	25.1
Short gestation and low birthweight nec (P07)		35	34	33	1	1
Rate		0.9	0.9	0.9	*	*
Sudden infant death syndrome (R95)		1,991	157	21	136	1,834
Rate		53.1	4.2	0.6	3.6	48.9
Maternal complications of pregnancy (P01)		26	24	23	1	1
Rate		0.7	0.7	0.6	*	*
Complications of placenta, cord, membranes (P02) ..		101	96	83	13	5
Rate		2.7	2.5	2.2	*	*
Respiratory distress of newborn (P22)		22	16	12	4	6
Rate		0.6	*	*	*	*
Accidents (unintentional injuries) (V01-X59)		769	77	19	57	692
Rate		20.5	2.0	*	1.5	18.5
Bacterial sepsis of newborn (P36)		117	110	54	56	7
Rate		3.1	2.9	1.4	1.5	*
Diseases of the circulatory system (I00-I99)		395	132	64	68	264
Rate		10.6	3.5	1.7	1.8	7.0
Intrauterine hypoxia and birth asphyxia (P20-P21)		322	294	193	101	28
Rate		8.6	7.8	5.2	2.7	0.8
All other causes		3,154	944	512	433	2,210
Rate		84.2	25.2	13.7	11.5	59.0
Not stated birthweight						
All causes		407	377	371	6	29
Rate		14,676.4	13,617.4	13,398.5	*	1,059.0
Congenital malformations (Q00-Q99)		31	26	25	1	5
Rate		1,108.8	926.2	890.1	*	*
Short gestation and low birthweight nec (P07)		175	175	174	1	-
Rate		6,315.2	6,315.2	6,278.9	*	-
Sudden infant death syndrome (R95)		3	1	1	-	2
Rate		*	*	*	-	*
Maternal complications of pregnancy (P01)		59	59	59	-	-
Rate		2,126.1	2,126.1	2,126.1	-	-
Complications of placenta, cord, membranes (P02) ..		39	39	39	-	-
Rate		1,390.4	1,390.4	1,390.4	-	-

See footnotes at end of table.

Documentation Table 5

Live births by birthweight and race of mother and infant deaths and infant mortality rates by age at death, birthweight, and race of mother for 10 major causes of infant death: United States, 2000 birth cohort data - Con.

[Infant deaths are weighted. Infant deaths are under 1 year. Neonatal deaths are under 28 days; early neonatal, 0-6 days; late neonatal, 7-27 days; and postneonatal, 28 days through 11 months. Rates are per 100,000 live births]

Cause of death, birthweight, and race of mother	Live births	Infant deaths	Total neonatal	Early neonatal	Late neonatal	Post-neonatal
All races ¹						
Not stated birthweight						
Respiratory distress of newborn (P22)		7	7	7	-	-
Rate		*	*	*	-	-
Accidents (unintentional injuries) (V01-X59)		2	1	1	-	1
Rate		*	*	*	-	*
Bacterial sepsis of newborn (P36)		5	5	3	2	-
Rate		*	*	*	*	-
Diseases of the circulatory system (I00-I99)		4	2	2	-	2
Rate		*	*	*	-	*
Intrauterine hypoxia and birth asphyxia (P20-P21)		8	7	6	1	1
Rate		*	*	*	*	*
All other causes		74	56	55	1	18
Rate		2,672.2	2,014.1	1,978.0	*	*
White						
All birthweights						
All causes		18,233	12,175	9,605	2,570	6,058
Rate		570.9	381.2	300.7	80.5	189.7
Congenital malformations (Q00-Q99)		4,451	3,271	2,509	763	1,180
Rate		139.4	102.4	78.5	23.9	36.9
Short gestation and low birthweight nec (P07)		2,383	2,338	2,260	78	45
Rate		74.6	73.2	70.8	2.4	1.4
Sudden infant death syndrome (R95)		1,616	129	18	111	1,487
Rate		50.6	4.0	*	3.5	46.6
Maternal complications of pregnancy (P01)		831	825	819	6	6
Rate		26.0	25.8	25.6	*	*
Complications of placenta, cord, membranes (P02) ..		713	705	680	25	8
Rate		22.3	22.1	21.3	0.8	*
Respiratory distress of newborn (P22)		621	574	458	117	46
Rate		19.4	18.0	14.3	3.7	1.5
Accidents (unintentional injuries) (V01-X59)		573	56	18	37	517
Rate		17.9	1.7	*	1.2	16.2
Bacterial sepsis of newborn (P36)		468	453	212	241	14
Rate		14.6	14.2	6.6	7.6	*
Diseases of the circulatory system (I00-I99)		447	172	96	76	276
Rate		14.0	5.4	3.0	2.4	8.6
Intrauterine hypoxia and birth asphyxia (P20-P21)		433	407	297	110	26
Rate		13.6	12.7	9.3	3.4	0.8
All other causes		5,698	3,245	2,239	1,007	2,452
Rate		178.4	101.6	70.1	31.5	76.8
Less than 2,500 grams						
All causes		11,227	9,338	7,851	1,487	1,889
Rate		5,359.4	4,457.7	3,747.8	709.9	901.7
Congenital malformations (Q00-Q99)		2,482	2,018	1,725	294	464
Rate		1,185.0	963.4	823.3	140.1	221.5

See footnotes at end of table.

Documentation Table 5

Live births by birthweight and race of mother and infant deaths and infant mortality rates by age at death, birthweight, and race of mother for 10 major causes of infant death: United States, 2000 birth cohort data - Con.

[Infant deaths are weighted. Infant deaths are under 1 year. Neonatal deaths are under 28 days; early neonatal, 0-6 days; late neonatal, 7-27 days; and postneonatal, 28 days through 11 months. Rates are per 100,000 live births]

Cause of death, birthweight, and race of mother	Live births	Infant deaths	Total neonatal	Early neonatal	Late neonatal	Post-neonatal
White						
Less than 2,500 grams						
Short gestation and low birthweight nec (P07)		2,265	2,221	2,145	76	44
Rate	1,081.3	1,060.4	1,024.1	36.3	20.9	
Sudden infant death syndrome (R95)		271	29	5	24	242
Rate	129.6	14.0	*	11.6	115.6	
Maternal complications of pregnancy (P01)		783	777	772	5	6
Rate	374.0	371.1	368.7	*	*	
Complications of placenta, cord, membranes (P02) ..		610	606	591	15	4
Rate	291.3	289.3	282.1	*	*	
Respiratory distress of newborn (P22)		604	561	446	115	42
Rate	288.1	267.9	213.1	54.8	20.2	
Accidents (unintentional injuries) (V01-X59)		61	7	4	3	54
Rate	29.0	*	*	*	25.6	
Bacterial sepsis of newborn (P36)		376	367	167	200	9
Rate	179.6	175.2	79.8	95.4	*	
Diseases of the circulatory system (I00-I99)		175	79	49	30	96
Rate	83.5	37.7	23.3	14.4	45.8	
Intrauterine hypoxia and birth asphyxia (P20-P21)		177	172	139	33	4
Rate	84.3	82.3	66.5	15.8	*	
All other causes		3,422	2,499	1,807	692	924
Rate	1,633.8	1,192.9	862.5	330.4	440.9	
2,500 grams or more						
All causes		6,754	2,608	1,530	1,078	4,146
Rate	226.5	87.5	51.3	36.2	139.0	
Congenital malformations (Q00-Q99)		1,946	1,233	764	469	713
Rate	65.3	41.4	25.6	15.7	23.9	
Short gestation and low birthweight nec (P07)		25	24	23	1	1
Rate	0.9	0.8	0.8	*	*	
Sudden infant death syndrome (R95)		1,342	99	12	86	1,243
Rate	45.0	3.3	*	2.9	41.7	
Maternal complications of pregnancy (P01)		12	12	11	1	-
Rate	*	*	*	*	-	
Complications of placenta, cord, membranes (P02) ..		77	73	63	10	4
Rate	2.6	2.5	2.1	*	*	
Respiratory distress of newborn (P22)		13	9	7	2	4
Rate	*	*	*	*	*	
Accidents (unintentional injuries) (V01-X59)		510	48	13	34	462
Rate	17.1	1.6	*	1.1	15.5	
Bacterial sepsis of newborn (P36)		88	83	44	39	5
Rate	3.0	2.8	1.5	1.3	*	
Diseases of the circulatory system (I00-I99)		269	92	46	46	178
Rate	9.0	3.1	1.5	1.5	6.0	
Intrauterine hypoxia and birth asphyxia (P20-P21)		250	229	153	76	21
Rate	8.4	7.7	5.1	2.5	0.7	

See footnotes at end of table.

Documentation Table 5

Live births by birthweight and race of mother and infant deaths and infant mortality rates by age at death, birthweight, and race of mother for 10 major causes of infant death: United States, 2000 birth cohort data - Con.

[Infant deaths are weighted. Infant deaths are under 1 year. Neonatal deaths are under 28 days; early neonatal, 0-6 days; late neonatal, 7-27 days; and postneonatal, 28 days through 11 months. Rates are per 100,000 live births]

Cause of death, birthweight, and race of mother	Live births	Infant deaths	Total neonatal	Early neonatal	Late neonatal	Post-neonatal
White						
2,500 grams or more						
All other causes		2,221	706	392	314	1,515
Rate		74.5	23.7	13.1	10.5	50.8
Not stated birthweight						
All causes		252	229	224	5	23
Rate		11,389.6	10,340.3	10,111.4	*	1,049.3
Congenital malformations (Q00-Q99)		23	20	20	-	3
Rate		1,020.6	*	*	-	*
Short gestation and low birthweight nec (P07)		93	93	92	1	-
Rate		4,187.0	4,187.0	4,141.5	*	-
Sudden infant death syndrome (R95)		3	1	1	-	2
Rate		*	*	*	-	*
Maternal complications of pregnancy (P01)		35	35	35	-	-
Rate		1,602.0	1,602.0	1,602.0	-	-
Complications of placenta, cord, membranes (P02) ..		25	25	25	-	-
Rate		1,142.7	1,142.7	1,142.7	-	-
Respiratory distress of newborn (P22)		4	4	4	-	-
Rate		*	*	*	-	-
Accidents (unintentional injuries) (V01-X59)		2	1	1	-	1
Rate		*	*	*	-	*
Bacterial sepsis of newborn (P36)		3	3	1	2	-
Rate		*	*	*	*	-
Diseases of the circulatory system (I00-I99)		3	1	1	-	2
Rate		*	*	*	-	*
Intrauterine hypoxia and birth asphyxia (P20-P21)		6	5	4	1	1
Rate		*	*	*	*	*
All other causes		55	41	40	1	14
Rate		2,473.1	1,833.4	1,788.3	*	*
Black						
All birthweights						
All causes		8,444	5,690	4,584	1,105	2,754
Rate		1,356.1	913.8	736.3	177.5	442.3
Congenital malformations (Q00-Q99)		1,022	683	513	170	339
Rate		164.1	109.7	82.4	27.3	54.4
Short gestation and low birthweight nec (P07)		1,833	1,799	1,756	43	33
Rate		294.4	289.0	282.0	7.0	5.4
Sudden infant death syndrome (R95)		770	67	10	57	702
Rate		123.6	10.8	*	9.2	112.8
Maternal complications of pregnancy (P01)		503	497	489	8	6
Rate		80.8	79.9	78.6	*	*
Complications of placenta, cord, membranes (P02) ..		284	278	267	11	6
Rate		45.7	44.7	42.9	*	*
Respiratory distress of newborn (P22)		342	322	262	60	20
Rate		55.0	51.7	42.1	9.7	3.2

See footnotes at end of table.

Documentation Table 5

Live births by birthweight and race of mother and infant deaths and infant mortality rates by age at death, birthweight, and race of mother for 10 major causes of infant death: United States, 2000 birth cohort data - Con.

[Infant deaths are weighted. Infant deaths are under 1 year. Neonatal deaths are under 28 days; early neonatal, 0-6 days; late neonatal, 7-27 days; and postneonatal, 28 days through 11 months. Rates are per 100,000 live births]

Cause of death, birthweight, and race of mother	Live births	Infant deaths	Total neonatal	Early neonatal	Late neonatal	Post-neonatal
Black						
All birthweights						
Accidents (unintentional injuries) (V01-X59)		271	31	10	21	240
Rate		43.5	5.0	*	3.4	38.5
Bacterial sepsis of newborn (P36)		275	260	108	153	15
Rate		44.2	41.8	17.3	24.5	*
Diseases of the circulatory system (I00-I99)		189	58	25	32	131
Rate		30.3	9.3	4.1	5.2	21.1
Intrauterine hypoxia and birth asphyxia (P20-P21)		146	138	98	40	8
Rate		23.5	22.2	15.8	6.5	*
All other causes		2,808	1,555	1,046	509	1,253
Rate		451.0	249.7	168.0	81.7	201.2
Less than 2,500 grams						
All causes		6,146	4,901	4,101	799	1,245
Rate		7,576.8	6,041.7	5,056.2	985.5	1,535.1
Congenital malformations (Q00-Q99)		591	430	347	83	160
Rate		728.2	530.7	427.8	102.9	197.5
Short gestation and low birthweight nec (P07)		1,755	1,721	1,678	43	33
Rate		2,163.3	2,122.1	2,068.7	53.4	41.2
Sudden infant death syndrome (R95)		201	14	3	11	187
Rate		248.4	*	*	*	231.1
Maternal complications of pregnancy (P01)		471	466	458	8	5
Rate		580.3	574.1	564.2	*	*
Complications of placenta, cord, membranes (P02) ..		256	250	242	8	6
Rate		315.6	308.1	298.1	*	*
Respiratory distress of newborn (P22)		332	314	255	59	18
Rate		409.5	387.2	314.0	73.2	*
Accidents (unintentional injuries) (V01-X59)		58	5	4	1	53
Rate		72.0	*	*	*	65.8
Bacterial sepsis of newborn (P36)		251	238	99	140	13
Rate		309.7	293.6	121.5	172.1	*
Diseases of the circulatory system (I00-I99)		85	27	13	14	57
Rate		104.5	33.7	*	*	70.8
Intrauterine hypoxia and birth asphyxia (P20-P21)		86	84	64	20	2
Rate		105.9	103.4	78.5	24.8	*
All other causes		2,060	1,351	940	411	709
Rate		2,539.4	1,665.4	1,158.3	507.1	874.0
2,500 grams or more						
All causes		2,170	664	358	306	1,506
Rate		400.8	122.7	66.2	56.5	278.2
Congenital malformations (Q00-Q99)		429	250	164	87	179
Rate		79.2	46.2	30.2	16.0	33.0
Short gestation and low birthweight nec (P07)		9	9	9	-	-
Rate		*	*	*	-	-
Sudden infant death syndrome (R95)		568	53	7	46	515
Rate		105.0	9.9	*	8.5	95.1

See footnotes at end of table.

Documentation Table 5

Live births by birthweight and race of mother and infant deaths and infant mortality rates by age at death, birthweight, and race of mother for 10 major causes of infant death: United States, 2000 birth cohort data - Con.

[Infant deaths are weighted. Infant deaths are under 1 year. Neonatal deaths are under 28 days; early neonatal, 0-6 days; late neonatal, 7-27 days; and postneonatal, 28 days through 11 months. Rates are per 100,000 live births]

Cause of death, birthweight, and race of mother	Live births	Infant deaths	Total neonatal	Early neonatal	Late neonatal	Post-neonatal
Black						
2,500 grams or more						
Maternal complications of pregnancy (P01)		11	10	10	-	1
Rate		*	*	*	-	*
Complications of placenta, cord, membranes (P02) ..		18	18	15	3	-
Rate		*	*	*	*	-
Respiratory distress of newborn (P22)		7	5	4	1	2
Rate		*	*	*	*	*
Accidents (unintentional injuries) (V01-X59)		213	26	6	20	187
Rate		39.3	4.8	*	3.7	34.5
Bacterial sepsis of newborn (P36)		22	20	7	13	2
Rate		4.1	3.7	*	*	*
Diseases of the circulatory system (I00-I99)		103	29	11	18	74
Rate		19.0	5.4	*	*	13.6
Intrauterine hypoxia and birth asphyxia (P20-P21)		59	52	32	20	6
Rate		10.8	9.7	6.0	3.7	*
All other causes		731	190	92	98	541
Rate		135.0	35.1	17.0	18.0	99.9
Not stated birthweight						
All causes		128	125	125	-	3
Rate		49,039.9	47,865.7	47,865.7	-	*
Congenital malformations (Q00-Q99)		2	2	2	-	-
Rate		*	*	*	-	-
Short gestation and low birthweight nec (P07)		69	69	69	-	-
Rate		26,423.4	26,423.4	26,423.4	-	-
Sudden infant death syndrome (R95)		-	-	-	-	-
Rate		-	-	-	-	-
Maternal complications of pregnancy (P01)		21	21	21	-	-
Rate		8,210.4	8,210.4	8,210.4	-	-
Complications of placenta, cord, membranes (P02) ..		10	10	10	-	-
Rate		*	*	*	-	-
Respiratory distress of newborn (P22)		3	3	3	-	-
Rate		*	*	*	-	-
Accidents (unintentional injuries) (V01-X59)		-	-	-	-	-
Rate		-	-	-	-	-
Bacterial sepsis of newborn (P36)		2	2	2	-	-
Rate		*	*	*	-	-
Diseases of the circulatory system (I00-I99)		1	1	1	-	-
Rate		*	*	*	-	-
Intrauterine hypoxia and birth asphyxia (P20-P21)		2	2	2	-	-
Rate		*	*	*	-	-
All other causes		17	14	14	-	3
Rate		*	*	*	-	*

* Figure does not meet standards of reliability or precision, see Technical notes.

- Quantity zero.

¹ Includes races other than white and black.

DOCUMENTATION TABLE 6

UNLINKED INFANT DEATHS BY RACE, AGE AT DEATH, AND STATE OF RESIDENCE:
 UNITED STATES, PUERTO RICO, VIRGIN ISLANDS, GUAM -- 2000 BIRTH COHORT DATA
 (INFANT DEATHS ARE UNDER 1 YEAR. NEONATAL DEATHS ARE UNDER 28 DAYS; EARLY NEONATAL,
 0-6 DAYS; LATE NEONATAL, 7-27 DAYS; AND POSTNEONATAL, 28 DAYS THROUGH 11 MONTHS)
 (DATA IN THIS TABLE IS FOR INFANT DEATHS IN 2000 THAT ARE NOT INCLUDED IN THE LINKED
 FILE BECAUSE THEY WERE NOT LINKED WITH THEIR CORRESPONDING BIRTH CERTIFICATES.
 SEE METHODOLOGY SECTION.

(RESIDENCE IS OF INFANT DECEDENT; RACE IS FROM DEATH CERTIFICATE.)

Area and Race of Child 1/ -----	Infant	Total NeoNatal	Early NeoNatal	Late NeoNatal	Post- NeoNatal
United States 2/.....	358	267	238	29	91
WHITE.....	240	176	153	23	64
BLACK.....	105	82	77	5	23
Alabama	-	-	-	-	-
WHITE.....	-	-	-	-	-
BLACK.....	-	-	-	-	-
Alaska	-	-	-	-	-
WHITE.....	-	-	-	-	-
BLACK.....	-	-	-	-	-
Arizona	5	4	3	1	1
WHITE.....	3	3	2	1	-
BLACK.....	-	-	-	-	-
Arkansas	1	-	-	-	1
WHITE.....	-	-	-	-	-
BLACK.....	1	-	-	-	1
California	56	48	44	4	8
WHITE.....	44	38	35	3	6
BLACK.....	10	9	9	-	1
Colorado	-	-	-	-	-
WHITE.....	-	-	-	-	-
BLACK.....	-	-	-	-	-
Connecticut	1	-	-	-	1
WHITE.....	-	-	-	-	-
BLACK.....	1	-	-	-	1
Delaware	1	1	1	-	-
WHITE.....	-	-	-	-	-
BLACK.....	1	1	1	-	-
District of Columbia	1	1	1	-	-
WHITE.....	-	-	-	-	-
BLACK.....	1	1	1	-	-
Florida	1	-	-	-	1
WHITE.....	1	-	-	-	1
BLACK.....	-	-	-	-	-
Georgia	-	-	-	-	-
WHITE.....	-	-	-	-	-
BLACK.....	-	-	-	-	-
Hawaii	4	2	2	-	2
WHITE.....	4	2	2	-	2
BLACK.....	-	-	-	-	-
Idaho	-	-	-	-	-
WHITE.....	-	-	-	-	-
BLACK.....	-	-	-	-	-
Illinois	12	6	6	-	6
WHITE.....	4	3	3	-	1
BLACK.....	8	3	3	-	5
Indiana	10	8	5	3	2
WHITE.....	9	7	4	3	2
BLACK.....	1	1	1	-	-
Iowa	-	-	-	-	-
WHITE.....	-	-	-	-	-
BLACK.....	-	-	-	-	-
Kansas	5	-	-	-	5
WHITE.....	4	-	-	-	4
BLACK.....	1	-	-	-	1

DOCUMENTATION TABLE 6

UNLINKED INFANT DEATHS BY RACE, AGE AT DEATH, AND STATE OF RESIDENCE:
 UNITED STATES, PUERTO RICO, VIRGIN ISLANDS, GUAM -- 2000 BIRTH COHORT DATA
 (INFANT DEATHS ARE UNDER 1 YEAR. NEONATAL DEATHS ARE UNDER 28 DAYS; EARLY NEONATAL,
 0-6 DAYS; LATE NEONATAL, 7-27 DAYS; AND POSTNEONATAL, 28 DAYS THROUGH 11 MONTHS)
 (DATA IN THIS TABLE IS FOR INFANT DEATHS IN 2000 THAT ARE NOT INCLUDED IN THE LINKED
 FILE BECAUSE THEY WERE NOT LINKED WITH THEIR CORRESPONDING BIRTH CERTIFICATES.
 SEE METHODOLOGY SECTION.

(RESIDENCE IS OF INFANT DECEDENT; RACE IS FROM DEATH CERTIFICATE.)

Area and Race of Child 1/ -----	Infant -----	Total NeoNatal -----	Early NeoNatal -----	Late NeoNatal -----	Post- NeoNatal -----
Kentucky	3	1	1	-	2
WHITE.....	3	1	1	-	2
BLACK.....	-	-	-	-	-
Louisiana	18	14	12	2	4
WHITE.....	4	3	3	-	1
BLACK.....	14	11	9	2	3
Maine	1	1	1	-	-
WHITE.....	1	1	1	-	-
BLACK.....	-	-	-	-	-
Maryland	10	6	6	-	4
WHITE.....	4	1	1	-	3
BLACK.....	6	5	5	-	1
Massachusetts	3	3	3	-	-
WHITE.....	1	1	1	-	-
BLACK.....	2	2	2	-	-
Michigan	1	-	-	-	1
WHITE.....	1	-	-	-	1
BLACK.....	-	-	-	-	-
Minnesota	1	1	1	-	-
WHITE.....	1	1	1	-	-
BLACK.....	-	-	-	-	-
Mississippi	-	-	-	-	-
WHITE.....	-	-	-	-	-
BLACK.....	-	-	-	-	-
Missouri	3	1	1	-	2
WHITE.....	3	1	1	-	2
BLACK.....	-	-	-	-	-
Montana	-	-	-	-	-
WHITE.....	-	-	-	-	-
BLACK.....	-	-	-	-	-
Nebraska	1	-	-	-	1
WHITE.....	1	-	-	-	1
BLACK.....	-	-	-	-	-
Nevada	7	2	1	1	5
WHITE.....	7	2	1	1	5
BLACK.....	-	-	-	-	-
New Hampshire	1	1	1	-	-
WHITE.....	1	1	1	-	-
BLACK.....	-	-	-	-	-
New Jersey	24	20	18	2	4
WHITE.....	14	12	12	-	2
BLACK.....	10	8	6	2	2
New Mexico	11	7	5	2	4
WHITE.....	8	6	4	2	2
BLACK.....	-	-	-	-	-
New York State	14	10	9	1	4
WHITE.....	10	7	6	1	3
BLACK.....	4	3	3	-	1
New York City	1	1	1	-	-
WHITE.....	1	1	1	-	-
BLACK.....	-	-	-	-	-
North Carolina	5	3	3	-	2
WHITE.....	3	2	2	-	1
BLACK.....	2	1	1	-	1

DOCUMENTATION TABLE 6

UNLINKED INFANT DEATHS BY RACE, AGE AT DEATH, AND STATE OF RESIDENCE:
 UNITED STATES, PUERTO RICO, VIRGIN ISLANDS, GUAM -- 2000 BIRTH COHORT DATA
 (INFANT DEATHS ARE UNDER 1 YEAR. NEONATAL DEATHS ARE UNDER 28 DAYS; EARLY NEONATAL,
 0-6 DAYS; LATE NEONATAL, 7-27 DAYS; AND POSTNEONATAL, 28 DAYS THROUGH 11 MONTHS)
 (DATA IN THIS TABLE IS FOR INFANT DEATHS IN 2000 THAT ARE NOT INCLUDED IN THE LINKED
 FILE BECAUSE THEY WERE NOT LINKED WITH THEIR CORRESPONDING BIRTH CERTIFICATES.
 SEE METHODOLOGY SECTION.

(RESIDENCE IS OF INFANT DECEDENT; RACE IS FROM DEATH CERTIFICATE.)

Area and Race of Child 1/ -----	Infant -----	Total NeoNatal -----	Early NeoNatal -----	Late NeoNatal -----	Post- NeoNatal -----
North Dakota	-	-	-	-	-
WHITE.....	-	-	-	-	-
BLACK.....	-	-	-	-	-
Ohio	50	47	41	6	3
WHITE.....	33	30	24	6	3
BLACK.....	15	15	15	-	-
Oklahoma	31	21	20	1	10
WHITE.....	20	11	10	1	9
BLACK.....	9	8	8	-	1
Oregon	-	-	-	-	-
WHITE.....	-	-	-	-	-
BLACK.....	-	-	-	-	-
Pennsylvania	3	1	-	1	2
WHITE.....	2	1	-	1	1
BLACK.....	1	-	-	-	1
Rhode Island	-	-	-	-	-
WHITE.....	-	-	-	-	-
BLACK.....	-	-	-	-	-
South Carolina	-	-	-	-	-
WHITE.....	-	-	-	-	-
BLACK.....	-	-	-	-	-
South Dakota	-	-	-	-	-
WHITE.....	-	-	-	-	-
BLACK.....	-	-	-	-	-
Tennessee	-	-	-	-	-
WHITE.....	-	-	-	-	-
BLACK.....	-	-	-	-	-
Texas	60	49	45	4	11
WHITE.....	44	36	33	3	8
BLACK.....	15	12	11	1	3
Utah	4	3	2	1	1
WHITE.....	4	3	2	1	1
BLACK.....	-	-	-	-	-
Vermont	-	-	-	-	-
WHITE.....	-	-	-	-	-
BLACK.....	-	-	-	-	-
Virginia	8	5	5	-	3
WHITE.....	4	2	2	-	2
BLACK.....	3	2	2	-	1
Washington	1	-	-	-	1
WHITE.....	1	-	-	-	1
BLACK.....	-	-	-	-	-
West Virginia	-	-	-	-	-
WHITE.....	-	-	-	-	-
BLACK.....	-	-	-	-	-
Wisconsin	-	-	-	-	-
WHITE.....	-	-	-	-	-
BLACK.....	-	-	-	-	-
Wyoming	-	-	-	-	-
WHITE.....	-	-	-	-	-
BLACK.....	-	-	-	-	-

DOCUMENTATION TABLE 6

UNLINKED INFANT DEATHS BY RACE, AGE AT DEATH, AND STATE OF RESIDENCE:
 UNITED STATES, PUERTO RICO, VIRGIN ISLANDS, GUAM -- 2000 BIRTH COHORT DATA
 (INFANT DEATHS ARE UNDER 1 YEAR. NEONATAL DEATHS ARE UNDER 28 DAYS; EARLY NEONATAL,
 0-6 DAYS; LATE NEONATAL, 7-27 DAYS; AND POSTNEONATAL, 28 DAYS THROUGH 11 MONTHS)
 (DATA IN THIS TABLE IS FOR INFANT DEATHS IN 2000 THAT ARE NOT INCLUDED IN THE LINKED
 FILE BECAUSE THEY WERE NOT LINKED WITH THEIR CORRESPONDING BIRTH CERTIFICATES.
 SEE METHODOLOGY SECTION.

RESIDENCE IS OF INFANT DECEDENT; RACE IS FROM DEATH CERTIFICATE.)

Area and Race of Child 1/ -----	Infant -----	Total NeoNatal -----	Early NeoNatal -----	Late NeoNatal -----	Post- NeoNatal -----
Puerto Rico 3/ WHITE.....	1 1	1 1	1 1	- -	- -
BLACK.....	-	-	-	-	-
Virgin Islands 3/ WHITE.....	5 1	3 1	2 -	1 1	2 -
BLACK.....	4	2	2	-	2
Guam 3/ WHITE.....	- -	- -	- -	- -	- -
BLACK.....	-	-	-	-	-

 /1 Totals for geographic areas include races other than white and black.
 /2 Excludes data for Foreign Residents, Puerto Rico, Virgin Islands, and Guam.
 /3 Data from the Puerto Rico, Virgin Islands, and Guam file.